



Photo by Ann Larson



Scentless Chamomile *Matricaria perforata* Mgmt Goal: **Suppression**

<p style="text-align: center;"><u>IDENTIFICATION</u></p> <ul style="list-style-type: none"> • Annual, reproducing by seed only; seed viable as soon as flower is formed. • 6-30" tall; bushy habit • Leaves fern-/feather-like; • Flower heads white with yellow center; 1/4–1 3/4" wide • Scentless — not the same variety as that used for tea 	<p style="text-align: center;"><u>CHEMICAL CONTROL</u></p> <p style="text-align: center;">CALIBRATE EQUIPMENT EVERY YEAR ALWAYS ADD NON-IONIC SURFACTANT ALWAYS FOLLOW THE HERBICIDE LABEL</p> <ul style="list-style-type: none"> • Roundup (glyphosate) <u>will also kill adjacent vegetation.</u> • Milestone (aminopyralid) <u>before flowering</u> <ul style="list-style-type: none"> ◦ Hand-Pump Sprayer: 1fl.oz. per 3 gal water ◦ All other sprayers: 4 6 to 6 fl.oz. per acre • Once flowers are present, herbicide will <u>not</u> prohibit seed spread.
<p style="text-align: center;"><u>IMPACTS</u></p> <ul style="list-style-type: none"> • Irritates mucus membranes of deer/elk. • Reduces/eliminates wildlife forage and habitat • Escaped Ornamental 	<p style="text-align: center;"><u>MECHANICAL CONTROL</u></p> <ul style="list-style-type: none"> • Once flowers are present, pull and bag flower heads to prevent seed spread. • Hand pull or dig when soil is moist. Make sure to pull up all roots.